

Elanco

DERBY DAYS



Get ready for the races with Elanco!



INCREASE SALES OF OUR INNOVATION
PRODUCTS AND GAIN NEW OCCUPANCIES!

PRODUCTS:

Zenrelia, Quattro, and CPMA

TIME:

May 1 - June 30, 2025

CLINICS CAN WIN 2 WAYS

For a new placement:

Zenrelia = \$2,400

Quattro = \$6,200

CPMA = \$1,320

For a re-order:

Zenrelia = \$1,200

Quattro = \$3,100

CPMA = \$1,320

PRIZES:

WIN

1 PRODUCT =
GARRETT'S POPCORN
SENT TO CLINIC

PLACE

2 PRODUCTS = 10
TUMBLERS

SHOW

3 PRODUCTS =
DERBY PARTY/HAT
BAR EXPERIENCE

Popcorn will be shipped within a week; Tumblers will be shipped out 2-4 weeks after promotion; Derby Party/Hat Bar will be live or virtual depending on clinic location. Questions? Contact Amy Lenzi at AMY.LENZI@elancoah.com.

INDICATIONS

Zenrelia is indicated for control of pruritus associated with allergic dermatitis and control of atopic dermatitis in dogs at least 12 months of age.

IMPORTANT SAFETY INFORMATION

Read the entire package insert before using this drug, including the Boxed Warning.
For Full prescribing information call 1-888-545-5973 or visit www.elancoh.com/us/zenrelia

WARNING: VACCINE-INDUCED DISEASE AND INADEQUATE IMMUNE RESPONSE TO VACCINES. Based on results of the vaccine response study, dogs receiving Zenrelia are at risk of fatal vaccine-induced disease and inadequate immune response to vaccines. Discontinue Zenrelia for at least 28 days to 3 months prior to vaccination and withhold Zenrelia for at least 28 days after vaccination. Dogs should be up to date on vaccinations prior to starting Zenrelia. Do not use in dogs less than 12 months old or dogs with a serious infection. Monitor dogs for infections because Zenrelia may increase susceptibility to opportunistic infections. Neoplastic conditions (benign and malignant) were observed during clinical studies. Consider the risks and benefits of treatment in dogs with a history of recurrence of these conditions. The most common adverse reactions were vomiting, diarrhea and lethargy. Zenrelia has not been evaluated in breeding, pregnant, or lactating dogs and concurrent use with glucocorticoids, cyclosporine, or other systemic immunosuppressive agents has not been tested. For full prescribing information see package insert.